| BUCYRUS COMMUNIT | | TATION STRATEGY RESULTS 2016 |
|---|--|--|
| Problem | Anticipated Long Term Outcomes (3 yrs) | Long Term Outcomes Results |
| Disability & Chronic Pain Arthritis/ <u>Rheumatism</u> Arthritis/ Rheumatism was pinpointed under Disability & Chronic Pain as being most needed by the community because 45.1% of people in Crawford County over the age of 50 suffer from arthritis/ rheumatism, 10% above the national average. | Decrease of arthritis sufferers from 45.1% to 44% in the 2016 community health needs assessment survey. | The 2016 CHNA was conducted by a different group and did not gather data for arthritis sufferers over the age of 50, but did gather data for sufferers over the age of 65. 59% of Crawford County Adults 60 years old and older suffer from arthritis. |
| Mental Health & Mental Disorders <u>Top Focus Group Concern</u> Resource Constraints/ Understaffed counseling services Crawford County in top 10 of 88 Ohio counties for adult suicides (2010 stats from American Foundation for Suicide Prevention) | Measured by a decrease in the Crawford County suicide rate as detailed by the American Society for Suicide Prevention and the CDC (statistics are between 1-2 years old) | UNKNOWN The only data available on the website of American Foundation for Suicide Prevention was at the state level. Ohio's rank in 2014 was 39 – There were 1,492 deaths by suicide in Ohio in 2014. |
| Substance Abuse Top Focus Group Concern Prevalence of Drug Use (especially Heroin) in Crawford County. | Decrease percentage of Crawford County residents who use drugs in the 2016 community health needs assessment survey. | AVITA HEALTH SYSTEM DID NOT ADDRESS THIS NEED, AS THERE IS A COUNTY WIDE OPIATE TASK FORCE. The 2016 community health needs assessment shows that 10% of adults use recreational drugs, including marijuana. |

| Galion Community Hospital CHNA Implementation Strategy Results 2016 | | | |
|--|--|--|--|
| Problem | Anticipated Long Term Outcomes (3 yrs) | Long Term Outcomes Results | |
| Nutrition & Overweight <u>Childhood Obesity</u> The percentage of children in Crawford County between the ages of 5-17 who are overweight is 43.2%, 13% higher than the national average of 30.7% | Measured by a decrease in the percentage of children in Crawford County between the ages of 5-17 who are overweight from 43.2% to 42% in the 2016 community health needs assessment survey. | Height and weight was not asked for in the 2016 CHNA survey, conducted by an agency different from the agency who conducted the 2012 CHNA survey. | |
| Tobacco Use <u>Current Smokers</u> 21.4% of Crawford County residents currently smoke, compared with national average of | Reduce percentage of current smokers in Crawford County to 21% on the next community health needs assessment survey. | The 2016 CHNA showed a reduction in the percentage of Crawford County smokers from 21% to 20% | |
| 16.6 [°] % <u>Smokers in the Home</u> 20.6% of Crawford County residents | Reduce percentage of smokers in the homes of Crawford County to 20% on the next community health needs assessment survey. | The 2016 CHNA showed an increase of smokers in the homes from 20% to 21%. | |
| reported that someone in the home smokes, versus 13.6% nationwide <u>Smoke in the Home with Children</u> 23.4% of Crawford County households with children have someone who smokes in the home, compared with 12.1% nationwide | Reduce percentage of people who smoke with children in the homes of Crawford County to 23% on the next community health needs assessment survey. | The 2016 CHNA showed that of the households with children, 18% had someone who smoked in the home. | |
| Heart Disease & Stroke <u>Heart Attack, Angina, Coronary</u> <u>Disease</u> 11.4% of people in Crawford County reported that they suffer from or have been diagnosed with heart disease, angina or heart attack, compared with a national average of 6.1% | Decrease of percentage of Crawford County residents diagnosed with stroke, heart disease, angina or heart attack from 11.4% to 10% in 2016 community health needs assessment survey. | Since the agency that conducted the 2016 CHNA report out the data in comparison to the BRFSS, heart attack and angina/coronary heart disease is reported out separately. They cannot combine the data to get the same percentage as reported before in the 2012 CHNA. The 2016 CHNA showed that 5% of Crawford County residents had angina and 5% had a heart attack, higher than the US average. It also showed that 1% of Crawford County residents had a stroke, which is higher than the US average. | |